

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
28 October 2004 (28.10.2004)

PCT

(10) International Publication Number  
**WO 2004/091624 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/496**,  
A61P 25/00, A61K 31/445

(21) International Application Number:  
PCT/EP2004/004126

(22) International Filing Date: 16 April 2004 (16.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0308968.7 17 April 2003 (17.04.2003) GB

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford Middlesex UB6 0NN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MELOTTO, Sergio** [IT/IT]; GlaxoSmithKline SpA, Via Alessandro Fleming 2, I-37100 Verona (IT). **CORSI, Mauro** [IT/IT]; GlaxoSmithKline SpA, Via Alessandro Fleming 2, I-37100 Verona (IT). **CARLETTI, Renzo** [IT/IT]; GlaxoSmithKline SpA, Via Alessandro Fleming 2, I-37100 Verona (IT).

(74) Agent: **GIDDINGS, Peter John**; GlaxoSmithKline, Corporate Intellectual Property, 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMBINATIONS COMPRISING PAROXETINE AND 2- (S) - (4-FLUORO-2-METHYL-PHENYL) -PIPERAZINE-1-CARBOXYLIC ACID [1- (R)- (3,5-BIS-TRIFLUORO-2-METHYL-PHENYL) -ETHYL]-METHYL AMIDE FOR TREATMENT OF DEPRESSION AND/OR ANXIETY

(57) Abstract: The present invention relates to therapeutic combinations comprising paroxetine or physiologically acceptable salts or solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, to pharmaceutical compositions containing said combinations and their use in the treatment of depression and /or anxiety.



WO 2004/091624 A1

COMBINATIONS COMPRISING PAROXETINE AND 2-(S)-(4-FLUORO-2-METHYL-PHENYL)-PIPERAZINE-1-CARBOXYLIC ACID [1-(R)-(3,5-BIS-TRIFLUORO-2-METHYL-PHENYL)-ETHYL]-METHYL AMIDE FOR TREATMENT OF DEPRESSION AND/OR ANXIETY

The present invention relates to therapeutic combinations comprising paroxetine or physiologically acceptable salts or solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, to pharmaceutical compositions containing said combinations and their use in the treatment of depression and /or anxiety.

Paroxetine ((-)- trans-4-(4'-fluorophenyl)3-(3'-4'-methylenedioxyphenoxy)methyl) piperidine) and its salts are commercially available and approved for use in humans for treatment and prophylaxis of, *inter alia*, anxiety, depression, obsessive compulsive disorder (OCD), premenstrual dysphoric disorder (PMDD) and panic disorders.

2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, which is described in WO01/25219, is a NK<sub>1</sub> receptor antagonist.

NK<sub>1</sub> receptor antagonists are known to be useful in the treatment of anxiety and depression, chemotherapy-induced nausea and vomiting and post-operative nausea and vomiting. Preclinical data suggest that NK<sub>1</sub> receptor antagonists may be useful in a variety of other disorders including pain, inflammatory diseases, allergic disorders, CNS disorders, skin disorders, cough and gastrointestinal disorders.

US 6117855 describes the use of a CNS-penetrant NK<sub>1</sub> receptor antagonist together with antidepressant or anti-anxiety drug for the manufacture of a medicament for the treatment or prevention of depression and/or anxiety.

There is however no specific disclosure of such combinations with paroxetine.

WO 01/25219 broadly teaches that the NK<sub>1</sub> receptor antagonists described therein may be administered in combination with a SSRI agent. However, there is no teaching concerning any synergistic effect of such combinations in the treatment of depression and /or anxiety.

It has now been found that, surprisingly, therapeutic compositions comprising a combination of paroxetine or physiologically acceptable salts or solvates thereof, for administration in combination with 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or

physiologically acceptable salts or solvates thereof to a human for the treatment of depression and/or anxiety, in which the dosage of the individual components are administered below the usual single therapeutic dosages, show surprising synergistic levels of efficacy for the treatment and/or prophylaxis of depression and/or anxiety.

5

In particular, it has now been found that by combining a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof a significantly greater antidepressant activity and/or anxiolytic activity than either of the two individual components taken alone is achieved .

It is a feature of this invention that the use of such a combination will provide one or more of the following effects: a more efficacious anti-depressive and/or anti-anxiety drug and/or a better tolerated drug treatment and/or a drug with a more rapid onset of the anti-depressive and/or anti-anxiety activity.

Furthermore, the synergistic effect of the combination of the present invention allows better management of any potential drug-related side effects.

According to one aspect of the invention, there is provided a combination comprising a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof.

25

When used in any of the contexts or aspects of the present invention a therapeutically non-effective dose refers to a dosage of each component of the combination which is lower than normally expected to produce effective therapeutic response when each component is administered alone.

30

When used in any of the contexts or aspects of the present invention, paroxetine or physiologically acceptable salts or solvates thereof, may be administered as the free base, or in the form of any physiologically acceptable salt thereof, including all hydrated or anhydrous forms and all polymorphic forms of such salts. In particular, references to paroxetine or physiologically acceptable salts or solvates thereof, include, without

35

limitation, paroxetine hydrochloride, paroxetine hydrochloride hemihydrate, paroxetine hydrochloride anhydrate, paroxetine mesylate and all polymorphic forms thereof.

Paroxetine is preferably used in the form of its hydrochloride hemihydrate salt.

5

Suitable pharmaceutically acceptable salts of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide include acid addition salts formed with pharmaceutically acceptable organic or inorganic acids, for example hydrochlorides, hydrobromides, sulphates, alkyl- or arylsulphonates (e.g. methanesulphonates or p-toluenesulphonates), phosphates, acetates, citrates, succinates, tartrates, fumarates and maleates.

10

Preferred physiologically acceptable salts of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide include hydrochloride, methanesulphonate, sulphate, p-toluenesulphonate.

15

2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide is preferably used in the form of its methanesulphonate salt.

20

According to the invention a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, may be in the range of 1 to 15 mg per day (measured as the free base), preferably in the range of 5 to 15 mg per day and most preferably in the range of 7 to 15 mg per day.

25

According to the invention a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof (measured as the free base) may be in the range of 1 to 10 mg per day, preferably in the range of 3.5 to 7.5 mg per day.

30

A combination according to the invention conveniently comprises paroxetine or physiologically acceptable salts or solvates thereof (measured as the free base), in an amount from 1 mg to 10 mg, more particularly in an amount from 3.5 mg to 7.5 mg, and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates

35

thereof, in an amount from 1 mg to 15 mg (measured as the free base) and particularly in an amount from 5 mg to 15 mg and more particularly in an amount from 7 to 15 mg.

5 A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount from 1 to 15 mg (measured as the free base).

10

A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount from 5 to 15 mg (measured as the free base).

15

A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount from 7 to 15 mg (measured as the free base).

20

25 A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount from 1 to 15 mg (measured as the free base).

30

A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically

35

acceptable salts or solvates thereof, in an amount from 5 to 15 mg (measured as the free base).

5 A preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount from 7 to 15 mg (measured as the free base).

10

A particularly preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof in an amount of 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount of 15 mg (measured as the free base).

15

Another particularly preferred combination according to the invention comprises paroxetine or physiologically salts or solvates thereof, in an amount of 3.75 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount of 15 mg (measured as the free base).

20

Another particularly preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount of 3.75 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount of 7.5 mg (measured as the free base).

25

30

Another particularly preferred combination according to the invention comprises paroxetine or physiologically acceptable salts or solvates thereof, in an amount of 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in an amount of 7.5 mg (measured as the free base).

35

The dose employed according to the present invention will of course depend on the method of administration, the age, the weight and condition of the patient.

- 5 The present invention thus provides a method for the treatment of depression and/or anxiety in a mammal including a human; which comprises treating said animal with a therapeutically effective amount of a combination comprising a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-  
10 carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof.

- In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said  
15 mammal with a therapeutically effective amount of a combination comprising a therapeutically non-effective dose of paroxetine and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

- 20 In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination comprising a therapeutically non-effective dose of paroxetine hydrochloride and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-  
25 (3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

- In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination comprising a  
30 therapeutically non-effective dose of paroxetine hydrochloride hemihydrate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

- 35 In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said

mammal with a therapeutically effective amount of a combination comprising a therapeutically non-effective dose of paroxetine hydrochloride anhydrate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination comprising a therapeutically non-effective dose of paroxetine mesylate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

Reference herein to treatment extends to prophylaxis as well as to treatment of established depression and/or anxiety symptoms.

As used herein, the term depression includes depressive mood episodes, depressive disorders, bipolar disorders, other mood, psychotic, adjustment disorders premenstrual and dysphoric disorder(PMDD). Thus, for example, depressive mood episodes include major depressive episodes and mixed episodes. Depressive disorders include Major Depressive Disorder (MDD) single or recurrent episode (with or without psychotic features, catatonic features, melancholic features, atypical features, anxious depression, or postpartum onset), dysthymic disorder (with early or late onset and with or without atypical features) and depressive disorder not otherwise specified. Bipolar disorders include bipolar I and II disorders, cyclothymic disorder and bipolar disorder not otherwise specified. Other mood, psychotic and adjustment disorders include neurotic depression; mood disorders due to general medical conditions including, but not limited to, myocardial infarction, diabetes, miscarriage, abortion, premenstrual dysphoric disorders(PMDD), dementia of the Alzheimer's type (with early or late onset) with depressed mood, vascular dementia with depressed mood; substance-induced mood disorders including, but not limited to, depression induced by alcohol, amphetamines, cocaine, hallucinogens, inhalants, opioids, phencyclidines, sedatives, hypnotics, anxiolytics and other substances; schizoaffective disorder of the depressed type; adjustment disorder with depressed mood; adjustment disorder with mixed anxiety and depressed mood.



As used herein, the term anxiety includes panic attacks, agoraphobia, anxiety disorders, adjustment disorders and separation anxiety disorder and premenstrual dysphoric disorder (PMDD). Thus, for example, anxiety disorders include panic disorder with or without agoraphobia, agoraphobia without a history of panic disorder, specific phobia, social phobia (social anxiety disorder), obsessive-compulsive disorder, Acute and posttraumatic stress disorders, generalised anxiety disorders, anxiety disorder due to a general medical condition, substance-induced anxiety disorder, anxiety disorder not otherwise specified and mixed anxiety-depression disorders. Adjustment disorders include adjustment disorder with anxiety and adjustment disorder with mixed anxiety and depressed mood.

The advantageous profile of anti-anxiety activity obtained by the administration of a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof with a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methylamide or a physiologically acceptable salt thereof can be demonstrated in the gerbil social interaction model, according to the method described by Cheeta et al. (Cheeta S. et al., 2001. Brain Research 915: 170-175).

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations, or sequentially. If there is sequential administration, the delay in administering the second and any subsequent active ingredient should not be such as to lose the benefit of a synergistic therapeutic effect of the combination of the active ingredients. It will also be understood that the compounds of the combination or the physiologically functional derivatives of any thereof, whether presented simultaneously or sequentially, may be administered individually or in multiples or in any combination thereof.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methylamide or a physiologically acceptable salt or solvates thereof in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine hydrochloride and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine hydrochloride hemihydrate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine hydrochloride anhydrate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

In a further preferred embodiment, the present invention provides the use of a therapeutically non-effective dose of paroxetine mesylate and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

The ratio of paroxetine or physiologically acceptable salts or solvates thereof to 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, in the combination according to the invention may be for example, from 1:15 to 10: 1 (measured by weight of the free bases), preferably from 1:4 to 4:1 (measured by weight of the free bases) and more preferably from 1:4 to 1:1 (measured by weight of the free bases).

The amount of a combination according to the invention required to be effective as an anti-depressive and/or anti-anxiety may, of course, vary and is ultimately at the discretion of the medical practitioner. The factors to be considered include the route of administration and nature of the formulation, the subject mammal's body weight, age and general condition and the nature and severity of the condition to be treated.

Unless otherwise indicated, all weights of active ingredients are calculated in terms of the drug *per se*. The desired dose may preferably be presented as one, two, three, four, five, six or more sub-doses administered at appropriate intervals throughout the day.

The components of the combination which may be referred to as active ingredients may be administered for therapy to an animal e.g. a mammal including a human in a conventional manner.

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. Pharmaceutical formulations according to the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. The carrier(s) must be acceptable in the sense of being compatible with the other ingredients of the formula and not deleterious to the recipient thereof. When the individual components of the combination are administered separately, they are generally each presented as a pharmaceutical formulation. The references hereinafter to formulations refer, unless otherwise stated, to formulations containing either the combination or a component thereof.

A combination of paroxetine or physiologically acceptable salts or solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-

phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, may conveniently be presented as a pharmaceutical formulation in a unitary dosage form.

5      Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package, usually a blister pack. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve  
10      patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment.

It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each formulation, containing within a package  
15      insert instructing the patient to the correct use of the invention is a desirable additional feature of this invention.

According to a further aspect of the invention provided is a multiple, for example, double or triple, pack comprising at least paroxetine or physiologically acceptable salts or  
20      solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, and an information insert containing directions on the use of the combination of the invention.

25      Formulations include those suitable for oral, rectal, nasal, topical (including transdermal, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods represent a further feature of the present invention and  
30      include the step of bringing into association the active ingredients with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

35      Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, caplets, cachets or tablets each containing a

predetermined amount of the active ingredients; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

5

A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as a powder or granules, optionally mixed with a binder (e.g. povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g. sodium starch glycollate, sodium croscarmellose cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface-active or dispersing agent. Molded tablets may be made by molding a mixture of the powdered compound moistened with an inert liquid diluent in a suitable machine. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredients therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach.

10

15

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredients in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredients in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredients in a suitable liquid carrier. Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or polyethylene glycols.

20

Topical administration may also be by means of a transdermal iontophoretic device.

Formulations suitable for vaginal administration may be presented as tablets, pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredients such carriers as are known in the art to be appropriate.

30

Pharmaceutical formulations suitable for rectal administration wherein the carrier is a solid are most preferably presented as unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be

35

conveniently formed by admixture of the active combination with the softened or melted carrier(s) followed by chilling and shaping in molds.

Formulations suitable for parenteral administration include aqueous and nonaqueous  
5 isotonic sterile injection solutions which may contain anti-oxidants, buffers, preservatives  
and solutes which render the formulation isotonic with the blood of the intended recipient;  
and aqueous and non-aqueous sterile suspensions which may include suspending agents  
and thickening agents; and liposomes or other microparticulate systems which are  
designed to target the compound to blood components or one or more organs. The  
10 formulations may be presented in unit-dose or multi-dose sealed containers, for example,  
ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring  
only the addition of the sterile liquid carrier, for example water for injection, immediately  
prior to use. Extemporaneous injection solutions and suspensions may be prepared from  
sterile powders, granules and tablets of the kind previously described.

15 It should be understood that in addition to the ingredients particularly mentioned above  
the formulations of this invention may include other agents conventional in the art having  
regard to the type of formulation in question, for example, those suitable for oral  
administration may include such further agents as sweeteners, thickeners and flavoring  
20 agents.

The pharmaceutical composition of the invention containing the two active ingredients  
may be prepared according to conventional techniques well known in the pharmaceutical  
industry. Thus, for example paroxetine or physiologically acceptable salts or solvates  
25 thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-  
trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates  
thereof, may be admixed together with suitable excipients such as those described above  
for the formulation of each of the active ingredients separately. Tablets may be prepared,  
for example by direct compression of such a mixture or using other conventional methods.  
30 Bilayer tablets may be prepared according to conventional procedure. Thus, for example,  
by separately compressing the two blends in a suitable tableting machine with two filling  
stations. Capsules may be prepared by filling the blend along with suitable excipients into  
gelatin capsules, using a suitable filling machine. Controlled release forms for oral or  
rectal administration may be formulated in a conventional manner associated with  
35 controlled release forms.

**Biological data:**

The advantageous profile of anti-anxiety activity obtained by the administration of a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof with a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-  
 5 amide or physiologically acceptable salts or solvates thereof, can be demonstrated in the gerbil social interaction model.

**Experiment**

10 Paroxetine hydrochloride hemihydrate (0.3 mg/kg p.o of the paroxetine measured as the free base.), 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate ( hereinafter compound A)(0.1 mg/kg i.p of the compound A measured as the free base) and a combination of  
 15 paroxetine (0.3 mg/kg p.o. of the paroxetine measured as the free base) and compound A (0.1 mg/kg i.p. of the compound A measured as the free base) were administered independently in mongolian gerbils to assess the effect on time spent in active social interactions.

The results obtained one hour after administration, expressed as a percentage variation of the time spent in active social interactions by each animal in respect to the value obtained  
 20 by treatment of control animals, are summarised in table 1.

Table 1

	<b>Paroxetine hydrochloride hemihydrate</b>	<b>Compound A</b>	<b>Combination of Paroxetine hydrochloride hemihydrate and Compound A</b>
<b>% variation</b>	<b>- 2%</b>	<b>- 8%</b>	<b>+ 56%</b>

The variation of the amount of time spent in active social interactions by each animal after treatment with a combination of paroxetine hydrochloride hemihydrate and compound A ,  
 25 is significantly greater than that expected from the therapeutic response of the components administered separately.

Thus, the above results provide evidence for a synergistic effect between 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-

ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof and paroxetine or physiologically acceptable salts or solvates thereof in a social stress assay.

5 Paroxetine as the free base or in the form of any physiologically acceptable salt thereof, including all hydrated or anhydrous forms and all polymorphic forms of such salts may be prepared by the method described in USP 4,007,196, EP-B-0223403, EP-B-0808314 and EP-B-0970955 which are incorporated herein by reference hereto.

10 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, may be prepared by the method described in WO 01/25219 which is incorporated herein by reference.

15 For co-administration, paroxetine or physiologically acceptable salts or solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, may be formulated in a conventional manner.

Thus, for example paroxetine or physiologically acceptable salts or solvates thereof may be formulated as described in USP 4,007,196, EP-B-0223403, EP-B-0808314 and EP-B-20 0970955 and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, may be formulated as described in WO 01/25219.

25 In a preferred aspect of the invention, paroxetine or physiologically acceptable salts or solvates thereof and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof, are formulated in a single pharmaceutical composition.

30 In order that this aspect of the invention may be more fully understood the following examples are given by way of illustration only.

In the following pharmaceutical formulation **Compound A** means 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof; **Compound B** 35 means paroxetine or physiologically acceptable salts or solvates thereof.



**Tablets**

Tablets may be prepared by the normal method such as direct compression or wet granulation.

The tablets may be film coated with a suitable film forming material such for example

5 Opadry using standard technique.

**Example 1****Direct compression****Tablets**

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)
Dibasic Calcium Phosphate	120.75 mg
Mg stearate	1.5 mg
Crospovidone	4.5 mg
Colloidal Silicon Dioxide:	0.75 mg

10

Compound A , Compound B, dibasic Calcium Phosphate, Crospovidone, Colloidal Silicon Dioxide and Magnesium stearate are mixed together and the resultant mix is compressed into tablets using suitable machine so as to provide tablets according to example 1.

15 **Example 2**

**Wet granulation**

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)
Microcrystalline cellulose	117.75 mg
Mg stearate	1.5 mg
Crospovidone	4.5 mg
Colloidal Silicon Dioxide	0.75 mg
Polyvinylpyrrolidone	3 mg

The compound A is mixed with Microcrystalline cellulose, Polyvinylpyrrolidone and Crospovidone then granulated with a suitable amount of water. After drying the granule, the compound B and Colloidal Silicon Dioxide are added to it and mixed for a suitable time. The resulting mixture was blended with Mg stearate and then compressed into tablets as described in Example 1.

### **Example 3**

#### **Wet granulation**

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)
Microcrystalline cellulose	117.75 mg
Mg stearate	1.5 mg
Crospovidone	4.5 mg
Colloidal Silicon Dioxide:	0.75 mg
Polyvinylpyrrolidone	3 mg

The compound B is mixed with Microcrystalline cellulose, Polyvinylpyrrolidone and Crospovidone then granulated with a suitable amount of water. After drying the granule, the compound A and Colloidal Silicon Dioxide are added to it and mixed for a suitable time. The resulting mixture was blended with Mg stearate and then compressed into tablets as described in Example 1.

### **Example 4**

#### **co Wet granulation**

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)
Microcrystalline cellulose	117.75 mg

Mg stearate	1.5 mg
Crospovidone	4.5 mg
Polyvinylpyrrolidone	3 mg

- The compound B and the compound A are mixed with Microcrystalline cellulose, Polyvinylpyrrolidone and Crospovidone then granulated with a suitable amount of water. After drying the granule, Mg stearate is added to it, blended and the resulting mixture
- 5 compressed into tablets as described in Example 1.

### **Example 5**

#### **Dry granulation**

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)
Microcrystalline cellulose	117.75 mg
Mg stearate	1.5 mg
Crospovidone	4.5 mg
Colloidal Silicon Dioxide:	1 mg
Polyvinylpyrrolidone	2 mg

- The compound B and the compound A are mixed with Microcrystalline cellulose, Mg stearate, Crospovidone, Colloidal Silicon Dioxide and Polyvinylpyrrolidone. The resulting mixture is compressed with flat faced punches so as to provide slugs which fall into a mill so as to obtain granular particles. The granule is then compressed into tablets as described in example 1.
- 10

### **Pellets**

### **Example 6**

#### Extrusion-Spheronization

<b>Compound A</b> (as methansulphonate salt)	15 mg (measured as the free base)
<b>Compound B</b> (as hydrochloride hemihydrate salt)	7.5 mg(measured as the free base)

Cellulose Spheres*	123 mg
Polyvinylpyrrolidone	4.5 mg

\* Microcrystalline cellulose (Avicel)

5 The compound B, after being mixed into the granulator chamber with microcrystalline cellulose, is wetted under agitation by spraying a suitable amount of water; the resulting wetted mass is extruded through a screen with proper dimensions so as to provide cylindrical extruded particles which are converted into pellets by the mechanical action of the rotating plate of a spheronizator. The pellets are dried and then encapsulated together with the compound A pellets produced applying the same process.

10 Alternatively, the compound B after being mixed into the granulator chamber with microcrystalline cellulose is wetted under agitation by spraying a suitable amount of water; the resulting mixture is agitated in order to let its particles growing up to pellets. The pellets are dried and then encapsulated together with the compound A pellets produced applying the same process.

15 Alternatively, a suitable amount of inert cellulose pellets are put in the fluid bed granulation chamber and set in motion introducing air at the bottom and then coated by spraying a water solution of the compound B. Pellets are dried and then encapsulated together with the compound A pellets produced applying the same process.

## CLAIMS

1. A combination comprising a therapeutically non-effective dose of paroxetine or physiologically acceptable salts or solvates thereof and a therapeutically non-effective dose of 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof.  
5
2. A combination as claimed in claim 1 wherein the paroxetine is present as its hydrochloride hemihydrate salt and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide is present as its methansulphonate salt.  
10
3. A combination as claimed in claim 1 or claim 2 comprising paroxetine or physiologically salts or solvates thereof in an amount from 1 to 10 mg (measured as the free base).  
15
4. A combination as claimed in any claims 1 to 3 comprising paroxetine or physiologically salts or solvates thereof in an amount from 3.5 to 7.5 mg (measured as the free base).  
20
5. A combination as claimed in any claims 1 to 4 comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 1 to 15 mg (measured as the free base).  
25
6. A combination as claimed in any claims 1 to 5 comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 5 to 15 mg (measured as the free base).  
30
7. A combination as claimed in any claims 1 to 6 comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 7 to 15 mg (measured as the free base).  
35

8. A combination as claimed in any claims 1 to 7 comprising paroxetine or physiologically salts or solvates thereof in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 1 to 15 mg (measured as the free base).
9. A combination as claimed in any claims 1 to 8 comprising paroxetine or physiologically salts or solvates thereof in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 5 to 15 mg (measured as the free base).
10. A combination as claimed in any claims 1 to 9 comprising paroxetine or physiologically salts or solvates thereof in an amount from 1 to 10 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 7 to 15 mg (measured as the free base).
11. A combination as claimed in any claims 1 to 10 comprising paroxetine or physiologically salts or solvates thereof in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 1 to 15 mg (measured as the free base).
12. A combination as claimed in any claims 1 to 11 comprising paroxetine or physiologically salts or solvates thereof in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 5 to 15 mg (measured as the free base).

13. A combination as claimed in any claims 1 to 12 comprising paroxetine or physiologically salts or solvates thereof in an amount from 3.5 to 7.5 mg (measured as the free base) and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically salts or solvates thereof in an amount from 7 to 15 mg (measured as the free base).
- 14 A combination as claimed in any claims 1 to 13 comprising paroxetine or physiologically acceptable salts or solvates thereof in an amount of 7.5 mg (measured as the free base) and comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof in an amount of 15 mg (measured as the free base).
- 15 15. A combination as claimed in any claims 1 to 14 comprising paroxetine or physiologically acceptable salts or solvates thereof in an amount of 3.75 mg (measured as the free base) and comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof in an amount of 15 mg (measured as the free base).
16. A combination as claimed in any claims 1 to 15 comprising paroxetine or physiologically acceptable salts or solvates thereof in an amount of 3.75 mg (measured as the free base) and comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof in an amount of 7.5 mg (measured as the free base).
17. A combination as claimed in any claims 1 to 16 comprising paroxetine or physiologically acceptable salts or solvates thereof in an amount of 7.5 mg (measured as the free base) and comprising 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or physiologically acceptable salts or solvates thereof in an amount of 7.5 mg (measured as the free base).

18. A combination according to any of claims 1 to 17 for use in the treatment and or prophylaxis of depression and /or anxiety.
- 5 19. A method for the treatment and/or prophylaxis of depression and/or anxiety in a mammal including a human, which comprises treating said animal with a therapeutically effective amount of a combination as claimed in any of claims 1 to 17.
- 10 20. A pharmaceutical formulation comprising a combination according to any of the claims 1 to 17 together with one or more pharmaceutically acceptable carriers or excipients.
- 15 21. The use of a combination according to any claims 1 to 17 in the manufacture of a therapeutically effective medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and /or anxiety.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/004126

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K31/496 A61P25/00 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	US 2003/144270 A1 (WROBLESKI MICHELLE LACI ET AL) 31 July 2003 (2003-07-31) page 8, paragraphs 173,174	1-21
Y	US 2002/123491 A1 (CHEN XIAO ET AL) 5 September 2002 (2002-09-05) page 3, paragraphs 36,37	1-21
Y	US 2003/064980 A1 (CHEN XIAO ET AL) 3 April 2003 (2003-04-03) page 2, paragraph 35 page 1, paragraph 4 page 3, paragraphs 38,39	1-21
Y	WO 02/10141 A (AHLIJANIAN MICHAEL KIRK ; SANNER MARK ALLEN (US); VILLALOBOS ANABELLA) 7 February 2002 (2002-02-07) page 42, lines 14,26	1-21
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

19 July 2004

Date of mailing of the international search report

27/07/2004

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Heller, D

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/004126

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/44200 A (CHEN XIAO ; PALIWAL SUNIL (US); XIAO DONG (US); BLYTHIN DAVID J (US);) 21 June 2001 (2001-06-21) page 5, paragraph 8-21 -----	1-21
Y	WO 01/25219 A (GLAXO GROUP LTD ; GIOVANNINI RICCARDO (IT); ST DENIS YVES (IT); ALVARO) 12 April 2001 (2001-04-12) cited in the application page 9, lines 3,4 page 14, lines 8-10 page 14, lines 22-26 -----	1-21
Y	WO 03/015784 A (JACK TORQUIL IAIN MACLEAN ; GLAXO GROUP LTD (GB); TRANQUILLINI MARIA E) 27 February 2003 (2003-02-27) page 2, line 15- page 5, lines 30-36 page 8, lines 8-11 -----	1-21

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2004/004126

### Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 19  
because they relate to subject matter not required to be searched by this Authority, namely:  
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/004126

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2003144270 A1	31-07-2003	WO 03042173 A1	22-05-2003
US 2002123491 A1	05-09-2002	US 2003064980 A1	03-04-2003
		AU 2097601 A	25-06-2001
		CA 2393672 A1	21-06-2001
		CN 1434805 T	06-08-2003
		EP 1237874 A2	11-09-2002
		HU 0301607 A2	29-12-2003
		JP 2003522739 T	29-07-2003
		WO 0144200 A2	21-06-2001
US 2003064980 A1	03-04-2003	US 2002123491 A1	05-09-2002
		AU 2097601 A	25-06-2001
		CA 2393672 A1	21-06-2001
		CN 1434805 T	06-08-2003
		EP 1237874 A2	11-09-2002
		HU 0301607 A2	29-12-2003
		JP 2003522739 T	29-07-2003
		WO 0144200 A2	21-06-2001
WO 0210141 A	07-02-2002	AU 7094401 A	13-02-2002
		BG 107469 A	30-09-2003
		BR 0112862 A	01-07-2003
		CA 2418115 A1	07-02-2002
		CN 1444567 T	24-09-2003
		CZ 20030225 A3	18-02-2004
		EP 1305295 A1	02-05-2003
		HR 20030048 A1	30-04-2003
		HU 0303069 A2	01-03-2004
		WO 0210141 A1	07-02-2002
		JP 2004505111 T	19-02-2004
		NO 20030472 A	27-03-2003
		SK 1042003 A3	04-05-2004
		US 2003083352 A1	01-05-2003
		US 2002119963 A1	29-08-2002
WO 0144200 A	21-06-2001	AU 2097601 A	25-06-2001
		CA 2393672 A1	21-06-2001
		CN 1434805 T	06-08-2003
		EP 1237874 A2	11-09-2002
		HU 0301607 A2	29-12-2003
		JP 2003522739 T	29-07-2003
		WO 0144200 A2	21-06-2001
		US 2003064980 A1	03-04-2003
		US 2002123491 A1	05-09-2002
WO 0125219 A	12-04-2001	AU 768780 B2	08-01-2004
		AU 7913900 A	10-05-2001
		BR 0014541 A	17-09-2002
		CA 2386515 A1	12-04-2001
		CN 1391564 T	15-01-2003
		CZ 20021166 A3	13-11-2002
		WO 0125219 A2	12-04-2001
		EP 1218359 A2	03-07-2002
		HU 0203136 A2	28-02-2003
		JP 2003511377 T	25-03-2003
		NO 20021637 A	06-06-2002
		NZ 518144 A	30-04-2004

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/004126

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0125219	A	TR 200200936 T2	22-07-2002
		US 2003028021 A1	06-02-2003
		US 2004048862 A1	11-03-2004
		ZA 200202589 A	03-07-2003
WO 03015784	A	27-02-2003	
		WO 03015784 A1	27-02-2003
		EP 1423117 A1	02-06-2004